

Computable structures of rank ω_1^{CK}

J. F. Knight and J. Millar

1 Introduction

We consider only computable languages and countable structures. In measuring complexity, we identify a structure with its atomic diagram. Makkai [9] produced an example of an arithmetical structure of Scott rank ω_1^{CK} . Here we obtain a computable structure of Scott rank ω_1^{CK} in two different ways. The first way is to start with Makkai's example and code it in a computable structure in a way that preserves the rank. We give a general coding procedure that transforms any hyperarithmetical structure $\mathcal{A}$ into a computable structure $\mathcal{A}^*$ such that the rank of $\mathcal{A}$ is ω_1^{CK} , or $\omega_1^{CK} + 1$, or $< \omega_1^{CK}$ iff the same is true of $\mathcal{A}^*$. The same coding procedure has been used recently in [7] for several further results.

The second way that we obtain a computable structure with Scott rank ω_1^{CK} is to examine Makkai's construction, and note that with a little more care, the resulting structure of Scott rank ω_1^{CK} can be made computable. We recall a construction, described nicely in [10], which associates trees with structures in such a way that a computable tree gives rise to a computable structure, and the tree has a path just in case the corresponding structure has a non-trivial automorphism. We show that if a computable tree is "thin", and has a path, but no hyperarithmetical one, then the resulting structure will have Scott rank ω_1^{CK} . Having done this, we use an idea of Sacks to obtain a tree with the desired properties.

The coding method is described in Section 2. In Sections 3 and 4, we re-work Makkai's construction. Section 3 gives the passage from trees to structures, and Section 4 constructs computable thin trees. In the remainder of the present section, we give a little background on Scott rank.

1.1 Scott rank

Scott [12] proved the following.

Theorem 1.1 (Scott Isomorphism Theorem) *If $\mathcal{A}$ is a countable structure for a countable language, there is an $L_{\omega_1, \omega}$ -sentence σ whose countable models are just the isomorphic copies of $\mathcal{A}$.*

In the proof of Theorem 1.1, Scott assigned countable ordinals to the tuples in a countable structure $\mathcal{A}$, and to the structure $\mathcal{A}$ itself. He started with the following family of equivalence relations $\equiv^\beta$ on tuples in $\mathcal{A}$.

1. $\bar{a} \equiv^0 \bar{b}$ if $\bar{a}$ and $\bar{b}$ satisfy the same quantifier-free formulas,
2. for $\beta > 0$, $\bar{a} \equiv^\beta \bar{b}$ if for all $\gamma < \beta$, for all $c \in \mathcal{A}$, there exists $d \in \mathcal{A}$ such that $\bar{a}, c \equiv^\gamma \bar{b}, d$ and for all $d \in \mathcal{A}$, there exists $c \in \mathcal{A}$ such that $\bar{a}, c \equiv^\gamma \bar{b}, d$.

Remark: We could split 2 into two parts.

- (a) For $\beta = \gamma + 1$, $\bar{a} \equiv^\beta \bar{b}$ if for all c there exists d , and for all d there exists c , such that $\bar{a}, c \equiv^\gamma \bar{b}, d$.
- (b) For limit β , $\bar{a} \equiv^\beta \bar{b}$ if for all $\gamma < \beta$, $\bar{a} \equiv^\gamma \bar{b}$.

If $\mathcal{A}$ is a countable structure, then for each tuple $\bar{a}$, there is some countable ordinal β such that all $\bar{b}$ satisfying $\bar{a} \equiv^\beta \bar{b}$ are in the orbit of $\bar{a}$. The *Scott rank* of the tuple $\bar{a}$, denoted by $SR(\bar{a})$, is the least such β . The *Scott rank* of the structure $\mathcal{A}$, denoted by $SR(\mathcal{A})$, is the least ordinal α such that for all tuples $\bar{a}$ in $\mathcal{A}$, $SR(\bar{a}) < \alpha$.

We are interested in computable structures and their ranks. We shall make use of *computable infinitary formulas* to state criteria for a computable structure to have various ranks. Roughly speaking, computable infinitary formulas are infinitary formulas in which the infinite disjunctions and conjunctions are over c.e. sets. For more information about these formulas, and for proofs of various facts stated below, see [2]. If $\mathcal{A}$ is a computable (or hyperarithmetical) structure, then for any tuple $\bar{a}$ in $\mathcal{A}$ and any computable ordinal β , there is a computable infinitary formula defining the $\equiv^\beta$ -class of $\bar{a}$. Using Barwise-Kreisel Compactness, we can show that the family of finite partial functions preserving $\equiv^{\omega_1^{CK}}$ has the back-and-forth property. It follows that if $\mathcal{A}$ is hyperarithmetical, then $SR(\mathcal{A}) \leq \omega_1^{CK} + 1$.

Proposition 1.2 *Suppose $\mathcal{A}$ is a hyperarithmetical structure.*

1. *If for some tuple $\bar{a}$, the orbit is not defined by any computable infinitary formula, then $SR(\mathcal{A}) = \omega_1^{CK} + 1$.*
2. *If for some computable ordinal α , the orbits of all tuples are defined by computable Π_α formulas, then $SR(\mathcal{A}) < \omega_1^{CK}$.*
3. *Otherwise, $SR(\mathcal{A}) = \omega_1^{CK}$.*

There are familiar examples of computable structures of Scott rank $\omega_1^{CK} + 1$. Harrison [8] showed that there is a computable ordering of type $\omega_1^{CK}(1 + \eta)$. In the Harrison ordering, the orbit of an element outside the well-ordered initial segment of type ω_1^{CK} is not defined by any computable infinitary formula, so the ordering has Scott rank $\omega_1^{CK} + 1$. There are related examples of Boolean algebras and Abelian p -groups which also have Scott rank $\omega_1^{CK} + 1$ [6].

Makkai [9] showed the following.

Theorem 1.3 (Makkai) *There exists an arithmetical structure $\mathcal{A}$ such that $SR(\mathcal{A}) = \omega_1^{CK}$.*

In the present paper, we give two different proofs of the following.

Theorem 1.4 *There exists a computable structure $\mathcal{A}$ such that $SR(\mathcal{A}) = \omega_1^{CK}$.*

2 Coding

The first proof of Theorem 1.4 takes Theorem 1.3 as given, and combines it with the result below.

Theorem 2.1 *Suppose $\mathcal{A}$ is a hyperarithmetical structure. Then there is a computable structure $\mathcal{A}^*$, with a copy of $\mathcal{A}$ definable in $\mathcal{A}^*$ by computable infinitary formulas, such that*

1. *If $\mathcal{A}$ has computable rank, then so does $\mathcal{A}^*$.*
2. *If $\mathcal{A}$ has rank ω_1^{CK} , then so does $\mathcal{A}^*$.*
3. *If $\mathcal{A}$ has rank $\omega_1^{CK} + 1$, then so does $\mathcal{A}^*$.*

In what follows, we first describe the structure $\mathcal{A}^*$. Then, we review some basic notions and results, before proving that the ranks of $\mathcal{A}$ and $\mathcal{A}^*$ are related as in Theorem 2.1.

2.1 Constructing $\mathcal{A}^*$

We suppose that $\mathcal{A}$ is a hyperarithmetical structure. We may suppose that the universe is computable and the language is relational. For each relation symbol P from the language of $\mathcal{A}$, and each tuple $\bar{a}$ from $\mathcal{A}$, having the arity of P , the pair $(P, \bar{a})$ corresponds to an infinite set of points in the structure $\mathcal{A}^*$. On this set, we put a copy of one of a pair of structures $\mathcal{C}_1, \mathcal{C}_2$, depending on whether $\mathcal{A} \models P\bar{a}$. The structures that we choose will be linear orderings.

We give a precise description of the structure $\mathcal{A}^*$, in terms of these linear orderings. There are unary relations L, L^*, A, U , and T , and binary relations E and $<$. The unary relations are disjoint, and their union is the universe of $\mathcal{A}^*$. The elements of L represent symbols in the language of $\mathcal{A}$. If $r \in L$ represents the i^{th} relation symbol, then r is connected via E to a cycle of length $i+1$ in L^* . The cycles are disjoint and exhaust L^* . The elements of A represent elements of $\mathcal{A}$.

For each n -tuple $a_0, \dots, a_{n-1}$ from $\mathcal{A}$, there is a corresponding n -tuple $u_0, \dots, u_{n-1}$ from U , where E connects u_i to a_i . In addition E takes u_i to u_{i+1} , and u_{n-1} to u_0 , forming a cycle. The cycles in U are all distinct, and they exhaust U . For each $r \in L$, representing an n -ary relation symbol R , and each n -tuple $\bar{a} = (a_0, \dots, a_{n-1})$ in $\mathcal{A}$, if u_0 is the element connected to a_0 in in the corresponding cycle in U , then u_0 and r are both connected via E to each

element in an infinite subset $T_{(R, \bar{a})}$ of T . The sets $T_{(R, \bar{a})}$ are disjoint and they exhaust T . There is one more binary relation on T , which is the union of the linear orderings on the separate sets $T_{(R, \bar{a})}$.

2.2 Some useful lemmas

We will choose the pair $\mathcal{C}_1, \mathcal{C}_2$ so that if $\mathcal{A}$ is Δ_α^0 , then we can make $\mathcal{A}^*$ computable. We use the lemma below, which was proved in [2] (see also [3]). We need some definitions. We write $(\mathcal{A}, \bar{a}) \leq_\beta (\mathcal{B}, \bar{b})$ if the Π_β formulas (of $L_{\omega_1, \omega}$) true of $\bar{a}$ in $\mathcal{A}$ are true of $\bar{b}$ in $\mathcal{B}$. If the tuples $\bar{a}$ and $\bar{b}$ are empty, we may write $\mathcal{A} \leq_\beta \mathcal{B}$. The relations $\leq_\beta$ are called the *standard back-and-forth relations*. Let α be a computable ordinal. A pair of structures $\mathcal{C}_1, \mathcal{C}_2$ is said to be α -friendly if the structures are computable and the relations, for $\beta < \alpha$, are uniformly c.e. For more about the standard back-and-forth relations, and the notion of α -friendliness, see [3].

Lemma 2.2 *Let α be a computable ordinal, and suppose $\mathcal{C}_1 \leq_\alpha \mathcal{C}_2$, where the pair $\{\mathcal{C}_1, \mathcal{C}_2\}$ is α -friendly. For any Π_α^0 set $S \subseteq \omega$, there is a uniformly computable sequence $(\mathcal{A}_n)_{n \in \omega}$ such that*

$$\mathcal{A}_n \cong \begin{cases} \mathcal{C}_2 & \text{if } n \in S \\ \mathcal{C}_1 & \text{otherwise} \end{cases}.$$

In [1], there is an analysis of the standard back-and-forth relations on computable ordinals that yields the next two lemmas.

Lemma 2.3 *For a computable ordinal β , $\omega^{\beta+1} \leq_{2\beta+1} \omega^\beta$ (so $\omega^\beta \leq_{2\beta} \omega^{\beta+1}$), and there is a computable $\Pi_{2\beta+1}^0$ sentence that is true in ω^β but not in $\omega^{\beta+1}$.*

Lemma 2.4 *For any computable ordinals α, β , there is an α -friendly pair of orderings of the types $\omega^\beta, \omega^{\beta+1}$.*

We need some further definitions and facts from [4], [5]. A computable (or X -computable) structure $\mathcal{A}$ is said to be *relatively intrinsically Δ_α^0 categorical* if for any copy $\mathcal{B}$ of $\mathcal{A}$, there is an isomorphism f from $\mathcal{A}$ onto $\mathcal{B}$ such that f is Δ_α^0 relative $\mathcal{B}$ (or X and $\mathcal{B}$). In [4] and [5], it is shown that a computable structure $\mathcal{A}$ is relatively intrinsically Δ_α^0 categorical iff it has a *formally Σ_α^0 Scott family*, where this is a Σ_α^0 set of computable Σ_α formulas, all with a fixed tuple of parameters, defining the orbits of all tuples in $\mathcal{A}$. If $\mathcal{A}$ has a formally Σ_α^0 Scott family with no parameters, then for any copy $\mathcal{A}$, then we get an isomorphism f which is $\Delta_\alpha^0(\mathcal{B})$ uniformly in $\mathcal{B}$. For an X -computable structure $\mathcal{A}$, the result is the same except that the Scott family is a $\Sigma_\alpha^0(X)$ set of X -computable Σ_α formulas.

A relation Q is *relatively intrinsically Σ_α^0* on a structure $\mathcal{A}$ if in all copies $\mathcal{B}$ of $\mathcal{A}$, the image of Q is Σ_α^0 relative to $\mathcal{B}$ and $\mathcal{A}$. In [4] and [5], it is shown that a relation is relatively Σ_α^0 on a computable structure $\mathcal{A}$ iff it is definable by a computable Σ_α formula with a finite tuple of parameters in $\mathcal{A}$.

Lemma 2.5 *Suppose $\mathcal{C}_1$ and $\mathcal{C}_2$ are uniformly relatively Δ_β^0 categorical, and each $\mathcal{C}_i$ satisfies a computable Π_β sentence that is not true in the other. Let $\mathcal{A}^*$ be as above.*

1. *If $\mathcal{A}$ is relatively intrinsically Δ_γ^0 -categorical, then $\mathcal{A}^*$ is relatively intrinsically $(\beta + \gamma)$ -categorical.*
2. *If $\mathcal{A}^*$ is relatively intrinsically Δ_γ^0 -categorical, then so is $\mathcal{A}$.*

Proof: For 1, suppose that $\mathcal{A}$ is relatively intrinsically Δ_γ^0 -categorical, let $\mathcal{B}^* \cong \mathcal{A}^*$, and let $\mathcal{B}$ be the copy of $\mathcal{A}$ that is naturally defined in $\mathcal{B}^*$, using computable Π_β formulas for the basic relations and their negations. Then $\mathcal{B}$ is $\Delta_\beta^0(\mathcal{B}^*)$, and there is an isomorphism from $\mathcal{A}$ to $\mathcal{B}$ that is $\Delta_\gamma^0(\mathcal{B})$, so it is $\Delta_{\beta+\gamma}^0(\mathcal{B}^*)$. We must extend this isomorphism to the rest of $\mathcal{A}^*$. For the part $T_{(R, \bar{a})}$, we locate the corresponding part of $\mathcal{B}^*$, and using $\Delta_\beta^0(\mathcal{B}^*)$, we can determine whether we are looking at a pair of copies of $\mathcal{C}_1$ or $\mathcal{C}_2$. For each pair of corresponding parts, we get an isomorphism that is $\Delta_\gamma^0(\mathcal{B}^*)$, uniformly. Putting the pieces together, we get an isomorphism from $\mathcal{A}^*$ onto $\mathcal{B}^*$ that is $\Delta_{\beta+\gamma}^0(\mathcal{B}^*)$.

For 2, suppose $\mathcal{A}^*$ is relatively intrinsically Δ_γ^0 -categorical, and let $\mathcal{B} \cong \mathcal{A}$. We may suppose that $\mathcal{B}$ has universe equal to the set of even numbers, and we can construct $\mathcal{B}^* \cong \mathcal{A}^*$ over the given $\mathcal{B}$, such that $\mathcal{B}^* \leq_T \mathcal{B}$. For an isomorphism from $\mathcal{A}^*$ onto $\mathcal{B}^*$ that is $\Delta_\gamma^0(\mathcal{B}^*)$, the restriction is an isomorphism from $\mathcal{A}$ onto $\mathcal{B}$ that is $\Delta_\gamma^0(\mathcal{B})$. Therefore, $\mathcal{A}$ is relatively intrinsically Δ_γ^0 -categorical.

Lemma 2.6 *Form $\mathcal{A}^*$ as above, using $\mathcal{C}_1$ and $\mathcal{C}_2$ which are relatively Δ_β^0 categorical, uniformly, where each satisfies a computable Π_β sentence that is not true in the other. Let Q be a relation on $\mathcal{A}$.*

1. *If Q is relatively intrinsically Σ_γ^0 on $\mathcal{A}$, then it is relatively intrinsically $\Sigma_{\beta+\gamma}^0$ on $\mathcal{A}^*$.*
2. *If Q is relatively intrinsically Σ_γ^0 on $\mathcal{A}^*$, then it is relatively intrinsically Σ_γ^0 on $\mathcal{A}$.*

The proof of Lemma 2.6 is like that for Lemma 2.5.

Suppose $\mathcal{A}$ is Δ_α^0 . We may assume that the universe is computable. Take β such that $\alpha \leq 2\beta + 1$, let $\mathcal{C}_1, \mathcal{C}_2$ be orderings of types $\omega^{\beta+1}, \omega^\beta$, respectively, and form $\mathcal{A}^*$ as described. By Lemmas 2.3, 2.4, and 2.5, we may take $\mathcal{A}^*$ to be computable. The orderings $\mathcal{C}_1, \mathcal{C}_2$ are both uniformly relatively $\Delta_{2\beta+2}^0$ categorical. Since there is a computable $\Pi_{2\beta+1}$ sentence true in ω^β but not in $\omega^{\beta+1}$, each $\mathcal{C}_i$ satisfies a computable $\Pi_{2\beta+2}$ sentence that is not true in the other. There is a $\Delta_{2\beta+2}^0$ copy of $\mathcal{A}$ definable in $\mathcal{A}^*$ —with an atomic formula defining the universe $\mathcal{A}$, and computable $\Pi_{2\beta+2}$ formulas defining the basic relations of $\mathcal{A}$ and their negations.

2.3 Proof of Theorem 2.1

We are ready to show that the Scott ranks of $\mathcal{A}$ and $\mathcal{A}^*$ are related as in Theorem 2.1. If the orbit of a tuple in one of the structures is intrinsically hyperarithmetical, then by a result of Soskov [13], it is definable by a computable infinitary formula without parameters (see also [6]). Suppose one of the structures has computable rank. Then that structure is relatively intrinsically hyperarithmetically categorical. By Lemma 2.6, the second structure is also relatively intrinsically hyperarithmetically categorical, so for some computable γ , there is a formally Σ_γ^0 Scott family, possibly with parameters. The orbit of the parameters is itself relatively intrinsically hyperarithmetical, so it is defined by a computable infinitary formula without parameters. Therefore, we can remove the parameters in the Scott family. From this, it follows that the second structure has computable Scott rank.

Next, suppose $\mathcal{A}$ has rank ω_1^{CK} . The orbits of all tuples in $\mathcal{A}$ are definable by computable infinitary formulas. Consider a tuple in $\mathcal{A}^*$. The part in $\mathcal{A}$ is described by a computable infinitary formula, and the part outside is as well. So, the orbits are all defined by computable infinitary formulas in $\mathcal{A}^*$. Suppose $\mathcal{A}^*$ has rank ω_1^{CK} . The orbits of all tuples in $\mathcal{A}^*$ are defined by computable infinitary formulas. In $\mathcal{A}$, the orbits are all relatively intrinsically hyperarithmetical, so they are defined by computable infinitary formulas, without parameters.

From what we have said, it follows that if one of the structures has rank $\omega_1^{CK} + 1$, then the other does as well. This completes the proof Theorem 2.1. Combining this with Makkai's Theorem (Theorem 1.3), we obtain a computable structure of rank ω_1^{CK} , as in Theorem 1.4.

3 From trees to structures

We have given one proof of Theorem 1.4, taking Theorem 1.3 as given. We shall re-work the proof of Theorem 1.3 to give another, self-contained, proof of Theorem 1.4. In this section, we reduce the proof to the construction of a computable tree with special properties. The construction is essentially due to Makkai. Morozov [10] used it to produce computable structures which share with the Harrison ordering the feature that there are non-trivial automorphisms, but not hyperarithmetical ones. We follow Morozov in the details.

Let T be a subtree of $\omega^{<\omega}$, and let T_n be the set of elements of T of length n . Let G_n be the set of finite subsets of T_n . For each n , G_n is an Abelian group under the operation of symmetric difference, denoted by Δ . The identity is $\emptyset$, and each element is its own inverse. In fact, G_n looks like a vector space over the 2-element field, with a basis consisting of $\{t\}$, for $t \in T_n$. For distinct $n > 0$, we replace $\emptyset$ by distinct elements id_n — G_0 consists of the single element $\emptyset = id_0$.

Let $G = \cup_n G_n$. The tree structure on T induces a tree structure on G , which is most easily described in terms of a *predecessor* function, which we will denote by p . If $a \in G_{n+1}$ represents a non-empty subset of T_{n+1} , say $a = \{t_1, \dots, t_k\}$, where t_i has predecessor t'_i (in the sense of T), then $p(a)$ is the sum (in G_n) of the

elements $\{t'_i\}$. We let $p(id_{n+1}) = id_n$. If $t \in T_n$ has two successors $t', t'' \in T_{n+1}$, then $p(\{t', t''\}) = id_n$. For each n , p acts as a homomorphism from G_{n+1} into G_n . Using the group structures, together with the tree structure, we define a family of unary operations f_a , for $a \in G$. If $a \in G_n$, then for $b \in G_m$, let $k = \min\{m, n\}$, and let a^* and b^* be the result of iterating $p(x)$ until the value is in G_k (i.e., $a^* = p^{(n-k)}(a)$ and $b^* = p^{(m-k)}(b)$). Then $f_a(b) = a^* \triangle b^*$ —this is the same as $f_b(a)$. Finally, we let $\mathcal{A}(T)$ be the structure $(G, (f_a)_{a \in G})$.

We have $a \in G_n$ iff $f_{id_n}(a) = a$ and for all $m < n$, $f_{id_m}(a) \neq a$. It follows that G_n is preserved under automorphisms of $\mathcal{A}(T)$. Similarly, p is preserved under automorphisms, since for $a \in G_{n+1}$ and $b \in G_n$, we have $p(a) = b$ iff $f_{id_n}(a) = b$. For any $a \in G_n$, $a = f_a(id_n)$. It follows that any automorphism of $\mathcal{A}(T)$ is determined by the images of the elements id_n .

Theorem 3.1 (Morozov) *For any computable tree $T \subseteq \omega^{<\omega}$, $\mathcal{A}(T)$ is a computable structure such that T has a path iff $\mathcal{A}(T)$ has a non-trivial automorphism. Moreover, if T has paths, then the Turing degrees of the paths are the same as the degrees of non-trivial automorphisms of $\mathcal{A}(T)$.*

Proof: Clearly, $\mathcal{A}(T)$ is computable in T . Suppose g is a non-trivial automorphism, and let $g_n = g(id_n)$. Since g is non-trivial, there is some n such that $g_n \neq id_n$. Take $t_n \in g_n$. For $m < n$, let t_m be the level m predecessor of t_n in T . By induction, for $m > n$, choose t_m to be a successor of t_{m-1} in g_m . These elements form a path through T . Conversely, if $(t_n)_{n \in \omega}$ is a path, we obtain a non-trivial automorphism g by mapping id_n to $\{t_n\}$, and then letting $g(a) = f_a(g(id_n))$, for $a \in G_n$.

To obtain a computable structure of rank ω_1^{CK} , we will produce a computable tree T with special features, and form the structure $\mathcal{A}(T)$. We need to develop an understanding of Scott rank in these structures.

3.1 Tree rank

We assign ranks to nodes of a tree T , and to T itself, as follows.

1. For $\sigma \in T$, $rk(\sigma) = 0$ if σ has no successors.
2. For $\alpha > 0$, $rk(\sigma) = \alpha$ if α is the first ordinal greater than $rk(\tau)$ for all successors τ of σ .

Note that $rk(\sigma) \geq \alpha + 1$ iff σ has a successor of rank $\geq \alpha$, and for limit α , $rk(\sigma) \geq \alpha$ iff $rk(\sigma) \geq \beta$ for all $\beta < \alpha$. We write $rk(\sigma) = \infty$ if σ does not have ordinal rank, where we consider that $\infty \geq \alpha$, for all ordinals α . We assign to the tree T the rank of the top node; i.e., $rk(T) = rk(\emptyset)$.

Lemma 3.2 *For all $\sigma \in T$, $rk(\sigma) = \infty$ iff σ lies on a path.*

We connect Scott rank in $\mathcal{A}(T) = (G, (f_a)_{a \in G})$ to tree rank in T . This will require several steps. First, for $a \in G$, we calculate the rank of a in the derived tree structure on G in terms of the ranks of elements of a in T .

Lemma 3.3 *Let $a \in G_n$, where $a \neq id_n$. Then $rk(a)$ is the minimum of $rk(t)$, for $t \in a$.*

Proof: Say $a = \{t_1, \dots, t_k\}$. We show, by induction on α , that $rk(a) \geq \alpha$ iff for all $t \in a$, $rk(t) \geq \alpha$. The statement is clearly true for $\alpha = 0$, and the clause for limit α is trivial. We consider $\alpha = \beta + 1$. First, suppose $rk(a) \geq \beta + 1$. Then a has a successor a^* of rank $\geq \beta$. By H.I., all $t \in a^*$ have rank $\geq \beta$. Now, $a^* = \{t_1^*, \dots, t_k^*\} \cup k$, where t_i^* is a successor of t_i of least rank in a^* , and k is a successor of id_n (if t_i has three successors in a^* , then t_i^* is one of least rank, and the other two are in k). Since t_i^* has rank $\geq \beta$, t_i has rank $\geq \beta + 1$. For the other direction, suppose that all t_i have rank $\geq \beta + 1$. For each i , let t_i^* be a successor of t_i of rank $\geq \beta$. Then $a^* = \{t_1^*, \dots, t_k^*\}$ is a successor of a , and by H.I., $rk(a^*) \geq \beta$. Therefore, $rk(a) \geq \beta + 1$.

From any single element $a \in G_n$, we can define any $b \in \cup_{m \leq n} G_m$, by the atomic formula $x = f_b(f_a(a))$. Using this, we see that if $a, a^* \in G_n$, then the function taking a to a^* extends in a unique way to an automorphism of the substructure $\cup_{m \leq n} G_m$,

Lemma 3.4 *Let $\bar{a}$ and $\bar{b}$ be tuples such that $\bar{a} \equiv^0 \bar{b}$. Suppose n is greatest such that some a_i is in G_n , and let $g = f_{a_i}(b_i)$. Then for all β , the following are equivalent:*

1. $\bar{a} \equiv^\beta \bar{b}$,
2. $\bar{a}, id_n \equiv^\beta \bar{b}, g$,
3. $id_n \equiv^\beta g$.

Proof: We proceed by induction. For $\beta = 0$, Statements 1, 2, and 3 are all true. The clause for limit β is trivial. We consider $\beta = \gamma + 1$. Supposing 1, we prove 2. For any c , there exists d , and for any d , there exists c , such that $\bar{a}, c \equiv^\gamma \bar{b}, d$. We suppose that c and d are in G_m for $m > n$ —this is the interesting case. Let $g' = f_c(d)$. By H.I., $\bar{a}, c, id_n, id_m \equiv^\gamma \bar{b}, d, g, g'$. Therefore, $\bar{a}, id_n, c \equiv^\gamma \bar{b}, g, c$. This gives 2. Statement 2 clearly implies Statement 3. Supposing 3, we prove 1. For any c , there exists d , and for any d , there exists c such that $id_n, c \equiv^\gamma a, d$. As above, we suppose that $c, d \in G_m$ for $m > n$ and let $g' = f_c(d)$. By H.I., we get $\bar{a}, c \equiv^\gamma \bar{b}, d$. This gives 1.

From Lemma 3.4, we get the following.

Lemma 3.5 *For any tuple $\bar{a}$, and n is greatest such that some $a_i \in G_n$, then $SR(\bar{a}) = SR(id_n)$.*

The Scott rank of id_n is calculated as follows.

Lemma 3.6 *For $a \in G_n$, $a \equiv^\beta id_n$ iff $rk(a) \geq \omega \cdot \beta$.*

Proof: We proceed by induction on β .

Case 1: $\beta = 0$.

We have a unique automorphism of the substructure $G_{\leq n} = \cup_{m \leq n} G_m$ taking id_n to a . It follows that for any atomic formula $\varphi(x)$, $\varphi(id_n)$ holds iff $\varphi(a)$ holds.

Case 2: $\beta = \gamma + 1$.

First, we suppose that $a \equiv^{\gamma+1} id_n$, and show that $rk(a) \geq \omega(\gamma+1)$. For each $k \in \omega$, there exists $c \in G_{n+k}$ such that $a, c \equiv^\gamma id_n, id_{n+k}$. Then $c \equiv^\gamma id_{n+k}$, and by H.I., $rk(c) \geq \omega\gamma$. Since $f_{id_n}(id_{n+k}) = id_n$, we must have $f_{id_n}(c) = a$, so c is a k^{th} successor of a . Therefore, $rk(a) \geq \omega\gamma + k$. Since this is true for all k , $rk(a) \geq \omega(\gamma+1)$.

Now, we suppose that $rk(a) \geq \omega(\gamma+1)$ and show that $a \equiv^{\gamma+1} id_n$. Suppose $b \in G_m$, where $m = n + k$ (this is the interesting case). Take $a^* \in G_m$ a k^{th} successor of a such that $rk(a^*) \geq \omega\gamma$. By H.I., $a^* \equiv^\gamma id_m$. Letting $c = f_b(a^*)$, we have $a^*, a, c \equiv^0 id_m, id_n, b$. Similarly, letting $c' = f_b(f_{a^*}(id_m))$, we have $a^*, a, b \equiv^0 id_m, id_n, c'$. By Lemma 3.4, we can replace $\equiv^0$ by $\equiv^\gamma$ in both cases. Then we get $a, c \equiv^\gamma id_n, b$ and $a, b \equiv^\gamma id_n, c'$. Therefore, $a \equiv^{\gamma+1} id_n$.

Case 3: β is a limit ordinal.

In this case, the statement holds by induction.

Having completed the proof of Lemma 3.6, we come to the main result of the section.

Theorem 3.7 *Suppose T is a computable tree such that for each n , there is a computable bound on the ordinal ranks of elements of T_n , but there is no computable bound on the set of ordinal ranks of elements of T . Then $\mathcal{A}(T)$ is computable, and $SR(\mathcal{A}(T)) = \omega_1^{CK}$.*

Proof: It is enough to show that for each n , $SR(id_n) < \omega_1^{CK}$, but there is no computable upper bound on these ranks. We have $SR(id_n) > \beta$ iff there exists a not in the orbit of id_n such that $rk(a) \geq \omega\beta$. For each n , there is a computable bound on the ordinal ranks of elements of G_n , say $\omega\beta_n$ is greater, but still computable. If $a \in G_n$ and $rk(a) \geq \omega\beta_n$, then a is in the orbit of id_n , so $SR(id_n) \leq \beta_n$. For each computable β , there exists n and $a \in G_n$ such that $rk(a)$ is an ordinal $\geq \omega\beta$. Then $a \equiv^\beta id_n$, but a is not in the orbit of id_n . This completes the proof of Theorem 3.7.

So, to produce a computable structure of rank ω_1^{CK} , it is enough to produce an appropriate computable tree, with a computable bound on the ordinal ranks at each level, but no computable bound over-all.

4 Thin trees

We define a special kind of tree satisfying the conditions of Theorem 3.7.

Definition: Let T be a subtree of $\omega^{<\omega}$, and let T_n be the set of nodes at level n . We say that T is *thin* if for each $n \geq 1$, the order type of the set of ordinal ranks of nodes in T_n is $\leq \omega \cdot n$.

Lemma 4.1 *If T is a computable thin tree, then for each n , there is a computable bound on the ordinal ranks of elements of T_n .*

Sacks suggested a way to produce computable thin trees of arbitrarily high computable rank.

Lemma 4.2 *For each computable ordinal α , there is a computable thin tree T , with no paths, such that $rk(T) \geq \alpha$.*

Proof: Let a be a notation for α (in Kleene's $\mathcal{O}$). We define a preliminary computable tree T_a , with labels, as follows. We give $\emptyset$ the label a . If σ has label 1 (representing 0), then σ has no successors. If σ has label $b = 2^c$, then σ has a single successor, with label c . If σ has label $3 \cdot 5^e$, then σ has an infinite family of successors, with labels b_n , where $\varphi_e(n) = b_n$. If $\sigma \in T$ has label b , where $|b| = \beta$, then β is the rank of σ in T_a . Then T_a has rank α . We replace T_a by a thin tree by slowing down the expansion of the limit notations.

We define the tree T , together with a computable function t from T_a to T . For σ at level 0 or 1 in T_a , $t(\sigma) = \sigma$. If $\sigma \in T$ has label of form 2^b , and σ' is the unique successor, with label b , then $t(\sigma)$ has a unique successor in T , and this is $t(\sigma')$. If $\sigma \in T_a$ has a label of form $3 \cdot 5^e$, where $\varphi_e(n) = b_n$, then we regard σ as “needing expansion”. We may delay expansion, in which case, $t(\sigma)$, will have a single successor, not in $ran(t)$. We may delay several times, giving $t(\sigma)$ a finite chain of successors. Say σ has successors σ_n with labels b_n . When we expand σ , we define $t(\sigma_n)$ to be a successor of the element σ' of length s in the chain below $t(\sigma)$.

At stage s in the construction, we have defined t mapping a finite subtree T_a^s of T_a into a finite subtree T^s of T , in which all terminal elements have length s or are t -images of elements with label $1 = |0|$. The subtree T_a^s may have some nodes σ which need expansion. If so, then at stage $s + 1$, we choose the first such σ , not previously acted on, and we begin the expansion process, letting t map the first successor of σ to a new successor of the level s extension of $t(\sigma)$. For other nodes $\sigma \in T_a^s$ such that $t(\sigma)$ has length s in T , we add a single successor, and we extend t , if appropriate. Finally, for each of the expansions begun earlier, we continue the expansion process. If we are expanding σ , then we extend t , mapping the next successor σ_n in $T_a - T_a^s$ to a new successor of $t(\sigma)$. In addition, we extend t further, and we add further nodes to T , so that the terminal nodes will all be at level $s + 1$ or be t -images of elements with label 1. The tree T obtained in this way is clearly computable.

It is easy to see that if $\sigma \in T$ has rank β , then $t(\sigma)$ has rank $\geq \beta$, so T has rank $\geq \alpha$. We can show, by induction on β , that if $\sigma \in T$ has rank β , then $t(\sigma)$ has rank $< \beta + \omega$. We must show that T is thin. We show by induction on $m \geq 1$ that the set of ordinal ranks of elements at level m is at most $\omega \cdot m$. At level 1, in T_a , there is either just one element or an infinite sequence σ_n with increasing ranks β_n . In the second case, note that for each n , $t(\sigma_n) = \sigma_n$, and the rank of σ_n in T is $\beta_n + k_n$, for some $k_n < \omega$. The sequence may not be strictly increasing, but for each n , there are only finitely many r such that $\beta_r + k_r < \beta_n + k_n$, so the set of ranks still has order type ω .

Supposing that the set of ranks at level m in T has order type at most $\omega \cdot m$, we show that the set of ranks at level $m + 1$ has order type at most $\omega \cdot (m + 1)$. There is at most one node at level m in T with infinitely many successors at level $m + 1$. All other nodes at level m have at most one successor. Say σ is the element of T_a chosen for expansion. Suppose σ has limit rank β , with successors σ_n of rank β_n . Let τ be the extension of $t(\sigma)$ at level m in T . The successors of τ are the elements $\tau_n = t(\sigma_n)$, where $\beta_n \leq rk(\tau_n) < rk(\beta)$. Therefore, τ has rank β . While $rk(\tau_n)$ need not be strictly increasing with n , since $rk(\tau_n)$ differs only finitely from β_n , for each n , there are only finitely many r such that $rk(\tau_r) < rk(\tau_n)$. For the elements at level m in T aside from τ , either the rank is 0, and there is no successor, or the rank is $\gamma + 1$, for some γ , and the unique successor has rank γ . So, at level $m + 1$ in T , the set of ranks is obtained by reducing the successor ranks that occur at level m by 1, and then adding an increasing sequence of ranks. The order type for level $m + 1$ is the result of taking m copies of ω , or an initial segment, and adding an ω sequence. The result has order type at most $\omega \cdot (m + 1)$.

Lemma 4.3 *There is a computable thin tree T such that $rk(T) = \infty$, but T has no hyperarithmetical path.*

Proof: We apply Barwise-Kreisel Compactness, using the previous lemma to satisfy the hypotheses. We have a Π_1^1 set Γ of computable infinitary sentences describing a model of KP in which the ordinals have an initial segment of type α for each computable ordinal α . We use a unary relation symbol T , and include sentences saying that it is a computable tree that does not have computable ordinal rank, but has no hyperarithmetical path. We add a binary relation symbol F , and include axioms saying that F maps elements of T to ordinals such that for $t \in T$ and α a computable ordinal, $F(t) = \alpha$ iff $rk(t) = \alpha$. Also, for each n , we add a binary relation symbol C_n , and we include axioms saying that C_n is an order-preserving function from $ran(F|T_n)$ onto an initial segment of $\omega \times n$.

Any Δ_1^1 subset of Γ is satisfied by taking the least admissible set as the model of KP , letting T be as in Lemma 4.2, of sufficiently high computable rank, with no path, taking F to be the rank function, and letting C_n be the function collapsing the ranks of elements of T_n to an initial segment of $\omega \cdot n$. Therefore, there is a model of the whole of Γ . In this model, the ordinals will not be well-founded. The function F will map tree elements that have ordinal

rank to their ranks, while mapping unranked elements to non-standard ordinals. The function C_n will map the ordinal ranks of elements of T_n that have ordinal rank to an initial segment of $\omega \cdot n$.

The fact that the tree in Lemma 4.3 does not have computable ordinal rank and has no hyperarithmetical path implies that there is no computable bound on the ordinal ranks of nodes. So, we get a tree T as in Theorem 3.7, and then $\mathcal{A}(T)$ is the structure we need for Theorem 1.4.

5 Categoricity

The Ryll-Nardjowski Theorem says that for an elementary first order theory T which is $\aleph_0$ categorical, there are only finitely many complete types in any fixed tuple of variables. Each type is principal, so there is a finite set of generating formulas. It is natural to wonder whether this generalizes. Suppose T is a computable infinitary theory which is $\aleph_0$ categorical. The types realized in models of T must be principal; i.e., for each type, there is a computable infinitary formula that generates it. However, for a given tuple of variables, there need not be a nice set of formulas generating the types. For the structures $\mathcal{A}(T)$ that we have been considering, the Ryll-Nardjowski Theorem fails badly.

Proposition 5.1 *If T is a computable thin tree which is unranked but has no hyperarithmetical path, then the computable infinitary theory of $\mathcal{A}(T)$ is $\aleph_0$ categorical. If Φ is a set of computable infinitary formulas defining the orbits of elements of $\mathcal{A}(T)$, then Φ cannot be hyperarithmetical, or even Π_1^1 .*

Proof: For each n , id_n has computable Scott rank, and the orbit is defined by a computable infinitary formula $\varphi_n(u)$. For any $a \in G_n$, the orbit of a is defined by $\exists u \varphi_n(u) \ \& \ x = f_a(u)$. There is no hyperarithmetical set of computable infinitary formulas defining the orbits of all elements, for then there would be a computable bound on the ranks. It follows that there can be no Π_1^1 set Φ of computable infinitary formulas defining the orbits of all elements, for then we could apply Barwise-Kreisel Compactness to produce a model of the computable infinitary theory of $\mathcal{A}(T)$ with an element a satisfying $\neg\varphi(x)$, for all $\varphi(x) \in \Phi$.

6 Problems

We close with some open problems.

The Harrison ordering $\mathcal{H}$, and the Harrison Boolean algebra the Harrison groups, have a strong computable approximation property.

Problem 1 *Is there a structure $\mathcal{A}$ of rank ω_1^{CK} such that for any Σ_1^1 set S , there is a uniformly computable sequence $(\mathcal{A}_n)_{n \in \omega}$ such that if $n \in S$, then $\mathcal{A} \cong \mathcal{A}_n$ and if $n \notin S$, then $\mathcal{A}_n$ has computable rank?*

Sacks and Young have further examples of structures of rank ω_1^{CK} with properties different from Makkai's example. Their examples are not hyperarithmetical but are "hyperarithmetically saturated", which means that the ordinals in the least admissible set over the structure are just the computable ordinals.

Problem 2 *Can we convert hyperarithmetically saturated structures of rank ω_1^{CK} into computable structures of this rank?*

References

- [1] Ash, C. J., "Categoricity in hyperarithmetical degrees", *Annals of Pure and Applied Logic*, vol. 34(1987), pp. 1–14.
- [2] Ash, C. J., and J. F. Knight, "Pairs of recursive structures", *Annals of Pure and Applied Logic*, vol. 46(1990), pp. 211–234.
- [3] Ash, C. J., and J. F. Knight, *Computable structures and the hyperarithmetical hierarchy*, Elsevier, 2000.
- [4] Ash, C. J., J. F. Knight, M. Mannasse, and T. Slaman, "Generic copies of countable structures", *Annals of Pure and Applied Logic*, vol. 42(1989), pp. 195–205.
- [5] Chisholm, J., "Effective model theory versus recursive model theory", *J. Symb. Logic*, vol. 55(1990), pp. 1168–1191.
- [6] Goncharov, S. S., V. S. Harizanov, J. F. Knight, and R. I. Shore, "Intrinsically Π_1^1 relations and paths through $\mathcal{O}$ ", submitted to *J. Symb. Logic*.
- [7] Goncharov, S. S., V. S. Harizanov, J. F. Knight, C. McCoy, R. G. Miller, and R. Solomon, "Enumerations in computable structure theory", submitted to *Annals of Pure and Applied Logic*.
- [8] Harrison, J., "Recursive pseudo well-orderings", *Transactions of the Amer. Math. Soc.*, vol. 131(1968), pp. 526–543,
- [9] Makkai, M., "An example concerning Scott heights", *J. Symb. Logic*, vol. 46(1981), pp. 301–318.
- [10] Morozov, A. S., "Groups of computable automorphisms", in *Handbook of Recursive Mathematics*, ed. by Yu. L. Ershov, et. al., 1998, Elsevier, pp. 311–345.
- [11] Sacks, G. E., *Higher Recursion Theory*, Berlin:Springer-Verlag, pp. 8–18.
- [12] Scott, D., "Logic with denumerably long formulas and finite strings of quantifiers", in *The Theory of Models*, ed. by J. Addison, L. Henkin, and A. Tarski, North-Holland, 1965, pp. 329–341.
- [13] Soskov, I. N., "Intrinsically hyperarithmetical sets", *Math. Logic Quarterly*, vol. 42(1996), pp. 469–480.